

INVESTIGATION OF PORCUPINE BEZOAR EXTRACT (HYSTRIX
BRACHYURAN) ON HeLa CELL COMBINED WITH ELECTROPORATION
METHOD

AZRA HAZWANIE BINTI AZIZULKARIM

A thesis submitted in
fulfillment of the requirement for the award of the
Degree of Master of Electrical Engineering

Faculty of Electrical and Electronic Engineering
Universiti Tun Hussein Onn Malaysia

JUNE 2019

ACKNOWLEDGEMENT

I would like to take this opportunity to express my greatest gratitude to Almighty Allah, for His help and guidance during this course of life and the moment of truth.

First and foremost, I would like to take this opportunity to express my appreciation and sincere gratitude to my supervisor, Associate Professor Ts. Dr. Muhammad Mahadi Bin Abdul Jamil and my co-supervisor Dr. Mohamad Nazib Bin Adon for their continuous support, encouragement and patient towards the completion of this Master research. I am thankful for their inspiring guidance, constructive criticism and advice throughout this project. I am honored and grateful to have an opportunity to work with them.

I too would like to dedicate this thesis to my beloved dad, Azizulkarim Bin Ibrahim for his tremendous patience, effort, support and encouragement. And for my late mom, Azizah Binti Nordin, this is for you.

My warmest appreciation to Dr. Salehudin bin Hamdan and Ms. Sayang binti Baba from Faculty of Bioscience and Medical Engineering, Universiti Teknologi Malaysia and to Dr Ridhwan bin Abd Wahab @ Abdul Wahab from the Department of Biomedical Science, Kulliyyah of Allied Health Sciences, International Islamic University Malaysia, for providing the required material and equipment for this research and also for their support and guidance.

Lastly, it has been a pleasure to be working with my best friend, Daing Noor Farhanah Binti Mohamad Ishak and I am most grateful for the help and support during my research. I would like to express my appreciation to UTHM, for the sponsorship via Contract Grant (Vote: U775) and finally thanks to everyone that may have contributed directly or indirectly for the accomplishment of this research.

ABSTRACT

Electroporation (EP) is a technique whereby the cell membrane is exposed to increase the intensity of electric field pulses that can lead to a variety of biophysical and biochemical responses. It is a molecular biology technique that stimulates pores through cell membrane, to increase the permeability of the cell, allowing chemicals, and drugs or Deoxyribonucleic acid (DNA) to be introduced into the cell. In this research, the HeLa cell behavior was investigated via EP. The purpose of this study is to investigate the best parameter of EP to be applied on HeLa cell treated with the best concentration of PBE which was 80.0ug/ml. Thus, the experiment is conducted for 48 hours and is initiated by identifying the best parameter for EP process. The range of electric field used is commenced between 100V/cm and 500V/cm with the increment of 100 V/cm interval and for the pulse duration from 100 μ s, 500 μ s, 1ms, 2ms and 5ms. Based on the parameter used, the best result obtained is when the parameter was set to 500 V/cm for the electric field and 100 μ s for pulse duration, where it inhibits the lowest proliferation rate percentage of HeLa cell. For the natural resource compound or extract which is available in Malaysia Porcupine Bezoar (*Hystrix Brachyuran*) were used. This extract has the potential as an anti-proliferation agent, which is useful in anti-cancer application. The experiment suggests that the best concentration for Porcupine Bezoar Extract (PBE) was at 80.0ug/ml. This combined method of EP and PBE revealed the process inhibited the proliferation of HeLa cell to 8.48%. In comparison to the study without this combined technique of EP and PBE, the inhibition process reached 55.35%. The study has successfully demonstrated the contribution towards a new method for anti-cancer treatment due to its minimal side effects compared to other conservative methods such as Electrochemotherapy (ECT).

ABSTRAK

Elektroporasi adalah satu teknik di mana sel membran terdedah kepada peningkatan intensity medan elektrik yang boleh membawa kepada pelbagai tindak balas biofizik dan biokimia. Ini adalah teknik biologi molekular untuk menghasilkan liang melalui sel membran, untuk meningkatkan kebolehtelapan sel, membolehkan bahan kimia, dan dadah atau DNA diperkenalkan ke dalam sel. Dalam kajian ini, ciri-ciri sel HeLa dikaji menggunakan teknik elektroporasi. Objektif kajian ini adalah untuk mengkaji parameter elektroporasi yang terbaik untuk digunakan pada sel HeLa yang juga dirawat menggunakan kepekatan terbaik PBE iaitu pada kepekatan 80.0 μ g/ml. Oleh itu, eksperimen telah dilakukan selama 48 jam dan bermula dengan menentukan parameter terbaik bagi proses elektroporasi. Julat voltan yang digunakan bermula dari 100V/cm, 200V/cm, 300V/cm, 400V/cm, dan 500V/cm, manakala untuk tempoh denyut yang digunakan bermula dari 100 μ s, 500 μ s, 1ms, 2ms dan 5ms. Berdasarkan semua parameter yang digunakan, parameter terbaik yang diperoleh adalah bila disetkan pada 500V/cm untuk julat voltan, dan 100 μ s untuk tempoh denyut yang mana dapat menghasilkan peratusan kadar proliferasi terendah bagi sel HeLa. Selain itu, sebatian atau ekstrak sumber asli yang terdapat di Malaysia dikenali sebagai *Hystrix Brachyuran* yang telah digunakan. Ekstrak ini berpotensi sebagai ejen anti-proliferasi yang berguna dalam proses anti-proliferasi dalam kajian anti-kanser. Dari eksperimen ini, kepekatan terbaik PBE adalah pada 80.0 μ g/ml. Kaedah gabungan antara EP dan PBE ini membuktikan bahawa proses untuk menghalang percambahan sel HeLa kepada 8.48%. Berbanding dengan kajian tanpa teknik gabungan EP dan PBE ini, proses pencambahan mencapai 55.35%. Oleh itu kajian menunjukkan potensi yang sangat besar sebagai sumbangan terhadap kaedah baru untuk rawatan anti-kanser yang mana mempunyai kesan sampingan yang minimal berbanding dengan kaedah konservatif yang lain seperti elektrokimoterapi (ECT).

TABLE OF CONTENTS

TITLE	i
ACKNOWLEDGEMENT	vi
ABSTRACT	vii
ABSTRAK	viii
TABLE OF CONTENTS	ix
LIST OF TABLES	xv
LIST OF FIGURES	xvi
LIST OF SYMBOL AND ABBREVIATIONS	xviii
LIST OF APPENDICES	xix
 CHAPTER 1 INTRODUCTION	 1
1.1 Background of Study	1
1.2 Problem Statement	2
1.3 Aim and Objectives	3
1.4 Scopes of Study	3
1.5 Thesis Structure	4
 CHAPTER 2 LITERATURE REVIEW	 6
2.1 The Cell	6
2.2 Composition of Cell	8
2.2.1 Nucleus	8
2.2.2 Cell Membrane	9
2.3 Hela Cell	10
2.3.1 Cell Attachment	11
2.3.2 Cell Proliferation	11
2.3.3 Cell Apoptosis	12
2.3.4 In-Vitro Method	13
2.4 Theory Of Electroporation (EP)	14
2.4.1 EP Types	15

2.4.1.1	Reversible Electroporation (RE)	15
2.4.1.2	Irreversible Electroporation (IRE)	15
2.4.2	Suitability Factors of the EP	16
2.4.2.1	Electric Field of EP	16
2.4.2.2	Pulse Duration of EP	17
2.4.2.3	Number of Pulses of EP	17
2.4.3	Application of EP	17
2.4.3.1	Electrochemotherapy (ECT)	17
2.4.3.1.1	Standard Electric Field Parameter for Electrochemotherapy (SEFPECT)	18
2.4.3.2	Electroinsertion	18
2.4.3.3	Electro-Sterilisation	19
2.4.4	Previous Study on EP Against Cancer	19
2.5	Porcupine Bezoar Extract (PBE)	20
2.5.1	History and Use of PBE	21
2.5.2	Previous Anti-Cancer Study by Using Natural Sources	22
CHAPTER 3	MATERIALS AND METHODS	24
3.1	Introduction	24
3.2	Materials for Hela Cell Line and Culture Conditions	26
3.2.1	Reagent for Hela Cell	26
3.2.2	Equipment	30
3.3	Methodology of the experiments	33
3.3.1	Protocol Culture of Hela Cell	33
3.3.2	Cell Sub-Culture Process	33
3.4	EP System and Technique	36
3.4.1	Square Wave Pulse Generator (ECM 830)	36
3.4.2	EP System	37
3.4.3	Experimental setup for EP	39
3.4.3.1	Experimental Setup for HeLa cell length measurement	40

3.4.3.2	Experimental setup for HeLa cell proliferation rate percentage by different pulse duration parameters	41
3.4.3.3	Experimental setup for HeLa cell proliferation rate percentage by different electric field parameters	42
3.5	TS100 Nikon Inverted Microscope	43
3.6	Dinocapture2.0 Software	43
3.6.1	Line Measurement Tool	44
3.7	Statistical Analysis	45
3.8	Porcupine Bezoar Extract (PBE) Preparation	45
3.9	Cell counting with Hemocytometer method	46
3.9.1	HeLaCell counting procedure for experiment treated with different concentrations of PBE	49
3.9.2	HeLa Cell counting procedure for experiment treated with both EP and PBE	50
3.10	Summary	51

CHAPTER 4 A PRELIMINARY EFFECT OF EP TECHNIQUE ON HELA CELL

4.1	Introduction	52
4.2	Materials and Methods	53
4.3	Results and Discussion	53
4.3.1	Result in Splitting Process for Hela Cell (Cell Culture Protocol)	53
4.3.2	Preliminary Results of EP on Hela Cell	55
4.3.3	Morphological Changes During EP	56
4.3	Summary	57

CHAPTER 5 SUITABILITY OF EP TECHNIQUE FOR HELA CELL

TOWARDS ANTI-PROLIFERATION APPLICATION 58

5.1	Introduction	58
5.2	Materials and Methods	59
5.3	Results and Discussion	60
5.3.1	Results of the EP Effect on Hela Cell With Different Pulse Duration after 48 hours	60
5.3.2	Discussions of the EP Effect on Hela Cell With Different Pulse Duration after 48 hours	61
5.3.3	Results of the EP effect on HeLa cell with Different Electric Field after 48 hours	63
5.3.4	Discussions of the EP effect on HeLa cell with Different Electric Field after 48 hours	64
5.3.5	Results of the comparison between HeLa cell induced with 500V/cm and 100 μ s of EP with normal HeLa cell after 48 hours	65
5.3.6	Discussions of the comparison between HeLa cell induced with 500 V/cm and 100 μ s of EP with normal HeLa cell after 48 hours	66
5.4	Summary	67

CHAPTER 6 PORCUPINE BEZOAR EXTRACT EFFECTS ON HELA CELL 68

6.1	Introduction	68
6.2	Materials and Methods	69
6.3	Results and Discussion	69
6.3.1	Results of the Effect of Different Concentrations Of PBE on the Proliferation Rate Percentage of HeLa Cell after 48 hours	69
6.3.2	Discussions of the effect of Different Concentrations of PBE on the Proliferation Rate Percentage of HeLa cell after 48 hours	71

6.3.3 Results of the Comparison Between Hela Cell Treated with 80.0µg/ml of PBE and Hela Cell Without Treatment with Natural Extract after 48 hours	72
6.3.4 Discussions of the Comparison Between HeLa Cell Treated with 80.0µg/ml of PBE and HeLa Cell Without Treatment with Natural Extract after 48 hours	73
6.4 Summary	74

CHAPTER 7 COMBINATION OF PORCUPINE BEZOAR

EXTRACT (PBE) WITH EP EFFECTS ON HELA CELL	75
7.1 Introduction	75
7.2 Materials and Methods	76
7.3 Results and Discussion	76
7.3.1 Results of the Comparison between HeLa Cell Proliferation Rate Percentage Treated with Both EP and PBE With Untreated HeLa Cell after 48 hours	76
7.3.2 Discussions of the Comparison Between HeLa Cell Proliferation Rate Percentage Treated With Both EP and PBE With Untreated HeLa Cell after 48 hours	78
7.3.3 Results of the Comparison of HeLa Cell Proliferation Rate Percentage Between Normal HeLa Cell as Control Group, HeLa Cell Induced With EP, HeLa Cell Treated With PBE and HeLa Cell With Both EP and PBE after 48 hours	79
7.3.4 Discussions of the Comparison of HeLa Cell Proliferation Rate Percentage Between Normal HeLa Cell as Control Group, HeLa Cell Induced With EP, HeLa Cell Treated With PBE and HeLa Cell With Both EP and PBE after 48 hours	82
7.4 Summary	82

CHAPTER 8 CONCLUSION AND RECOMMENDATION	84
8.1. Conclusion	84
8.2. Recommendation for Future Work	85
REFERENCES	86
APPENDIX	94
LIST OF PUBLICATIONS AND COPYRIGHTS	135
VITA	137



LIST OF TABLES

3.1	Type of EP Parameter	36
5.1	Percentage mean, Standard Deviation and Standard Error of HeLa Cell Proliferation Rate Percentage for Different Pulse Duration Parameter after 48 Hours	60
5.2	Percentage mean, Standard Deviation and Standard Error of HeLa Cell Proliferation Rate Percentage for Different Electric Field Parameter after 48 Hours	63
5.3	Percentage Mean, Standard Deviation and Standard Error Between HeLa Cell Proliferation Rate Percentage Induced With EP and Without EP after 48 Hours	65
6.1	Percentage Mean, Standard Deviation and Standard Error of HeLa Cell Proliferation Rate Percentage for Different Concentrations of PBE after 48 Hours	70
6.2	Percentage Mean, Standard Deviation and Standard Error Between HeLa Cell Proliferation Rate Percentage Treated With PBE Versus Untreated HeLa Cell after 48 hours	72
7.1	Percentage Mean, Standard Deviation and Standard Error Between HeLa Cell Proliferation Rate Percentage Treated With Both EP and PBE With Untreated HeLa Cell after 48 hours	77
7.2	Percentage Mean, Standard Deviation and Standard Error Between Normal HeLa Cell as Control Group, HeLa Cell Induced With EP, HeLa Cell Treated With PBE and HeLa Cell With Both EP and PBE after 48 hours	80

LIST OF FIGURES

2.1	A Diagram of A Prokaryotic Cell	7
2.2	Eukaryotic Cell	7
2.3	Illustration of Nucleus	8
2.4	Illustration of Cell Membrane	9
2.5	Cervical Cancer Cell (Hela Cell)	10
2.6	Illustration of Cell Apoptosis	13
2.7	The Difference Between RE AndIRE	16
2.8	Image of Landak Raya or Also Known As Landak Borneo By Locals	21
3.1	The Flowchart of The General Methodology of The Research	25
3.2	RPMI 1640 Media	27
3.3	Fetal Bovine Serum (FBS)	27
3.4	TrypLe Express Solution	28
3.5	Penicillin/Streptomycin	29
3.6	Phosphate Buffer Saline (PBS)	29
3.7	Trypan Blue	30
3.8	Incubator	31
3.9	Water Bath	31
3.10	Centrifuge Machine	32
3.11	Biosafety Cabinet	32
3.12	ECM 830 Electroporator	33
3.13	Comparison Between The Hela Cell Condition (A) Before Detached And (B) After Trypinized With Tryple express And Start To Detached From The Surface	35
3.14	ECM830 Square Wave Pulse Generator.	37
3.15	EP System	38

3.16	Image of Cuvette That Have An Electrode Gap of 4mm with Electric Field Stimulation	38
3.17	The Image of Nikon TS100 Inverted Microscope Equipped with Dino Camera And Dinocapture2.0 Software	39
3.18	Image of Line Measurement Equipped with Dinocapture2.0 Software	45
3.19	The Image Illustrate The Procedure of PBE Preparation	46
3.20	(a-b) A Hemocytometer schematic layout	48
3.21	(a-c) Hemocytometer method to counting the number of HeLa cell	
4.1	The Image of HeLa Cell After Reach 80-90% Confluences in 25cm ² Flask. (Scale Bar = 20µm).	53
4.2	HeLa cell proliferation rate in 48 hours. (Scale bar = 20µm)	54
4.3	HeLa cell cultured on culture flask for 48 hours (Scale bar =20µm)	55
4.4	Comparison sized of HeLa cell a) without EP and b) with EP (Scale bar = 20µm).	56
5.1	The Relationship between Electric Field and Pulse Duration. Illustration Adapted From (Dower <i>Et Al.</i> , 1988).	59
5.2	Graph of HeLa cell Proliferation Rate Percentage for Different Pulse Duration Parameter after 48 hours	61
5.3	Graph of HeLa cell Proliferation Rate Percentage for Different Electric Field Parameter after 48 hours	63
5.4	Graph of comparison between proliferation rate percentage of HeLa cell treated with EP and without EP after 48 hours	66
6.1	Graph of HeLa cell ProliferationRate Percentage for Different Concentrations of PBE after 48 hours	70
6.2	Graph of comparison between proliferation rate percentage of HeLa cell treated with PBE and without PBE after 48 hours	72
7.1	Graph of HeLa cell proliferation rate percentage treated with both PBEwith EP technique versus untreated HeLa cell after 48 Hours	77
7.2	Graph of comparison of HeLa cell proliferation rate percentage between normal HeLa cell as control group, HeLa cell induced with EP, HeLa cell treated with PBE and HeLa cell with both EP and PBEafter 48 hours	81

LIST OF SYMBOL AND ABBREVIATIONS

<i>DNA</i>	-	Deoxyribonucleic acid
<i>ECM 830</i>	-	Square Wave Pulse Generator
<i>ECT</i>	-	Electrochemotherapy
<i>EP</i>	-	Electroporation
<i>FBS</i>	-	Fetal Bovine Serum
<i>HeLa</i>	-	Cervical Cancer Cell
<i>IRE</i>	-	Irreversible Electroporation
<i>PBE</i>	-	Porcupine Bezoar Extract
<i>PBS</i>	-	Phosphate Buffer Saline (PBS)
<i>PEF</i>	-	Pulse Electric Field
<i>RE</i>	-	Reversible Electroporation
<i>RPMI 1640</i>	-	Roswell Park Memorial Institute Medium
<i>SEFPECT</i>	-	Standard Electric Field Parameters for Electrochemotherapy
<i>TPC</i>	-	Total phenolic content
<i>TFC</i>	-	Total flavanoid content

LIST OF APPENDICES

APPENDIX	TITTLE	PAGE
A	Published journals and conference papers	94
B	Safety Data Sheet	97



PTTA UTHM
PERPUSTAKAAN TUNKU TUN AMINAH

CHAPTER 1

INTRODUCTION

1.1 Background of study

Cancer is one of the leading factors of human casualty which is approximate at 8.2 million globally per year and most likely to increase to 13 million worldwide per year until 2030 (Mehta *et al.*, 2014). Moreover, oncology has become the biggest therapeutic area in the pharmaceutical sector in terms of the number of project clinical trials on research and development spending (Moffat *et al.*, 2014).

The past decade has been an escalation in cancer biology research with many new promising disease targets. However, the capability to translate these advances into therapies are poor, with a failure rate of about 90% (Adams *et al.*, 2012). Budget reduction on studies and development by pharmaceutical companies and expenditure increment of bringing a new molecular entity to market, the anti-drug advancement had becomes a highly complex problem (Adams *et al.*, 2012). Moreover, advances in cancer cure have proved in high number of long – term cancer survivors whom most of them were left to experience the side effect from their treatments such as hair loss, joint pain and nausea. One of the contributing factor includes the drugs intake during treatment. For over 50 years, the investigation of anti-cancer drugs has been directed by the fact that tumour cell replicates more expeditiously than normal cell and that DNA is the most essential element in cell division. DNA is a common therapeutic

target of anti-cancer drugs, and most of the currently available drugs may cause DNA damages, interrupts cell division and finally causes cell death (Bailón-Moscoso *et al.*, 2014).

Studies of anti-cancer, by using animal extract as a chemotherapeutic agent are investigated rapidly. The use of Porcupine Bezoar Extract (PBE) in this research will be explored of its capability to inhibit the proliferation rate percentage of HeLa cell through Electroporation (EP). This is accomplished by using EP with animal extract. Bioactive extract from porcupine bezoar which is also known as the “prince of antidotes” (Barroso MD and Duffin CJ, 2013), that has the capabilities of anti-proliferation and also anti-angiogenesis properties for cancer growth is employed. Porcupine bezoar or porcupine stone is extracted from porcupine rodents with a coat or sharp quills, which normally will defend and camouflage themselves from predators like snakes or tigers. Since conventional treatment often cause certain side effect to cancer survivors, PBE will be explored to whether it reduces or inhibit the proliferation rate percentage of HeLa cell through EP technique. This could serve as an alternative method for cancer treatment as it originates naturally and, with minimal or no side effects on its application.

1.2 Problem statement

Presently, cancer is known to be the leading cause and factor of death throughout the world. It is one of the major threats to public health in both developed and developing nations. In developed countries, cancer is the second most common cause of death (Moffat *et al.*, 2014). Studies show that cancer is caused by both external and internal factors. The external factors are tobacco, consumption, radiations, chemicals, and infectious organism whilst internal factors include inherited mutations, hormones and also immune condition related to sedentary lifestyles likes smoking, poor diet and inactive physical activities (Mehta *et al.*, 2014).

Various treatments are provided in order to help cancer patients, for instance chemotherapy, immune treatment, radiation therapy, hyperthermia, hormone therapy and many more. Consequently, advances in cancer therapy will lead and result in increasing numbers of long-term survivors who are left to themselves to handle the

consequences of their treatments. Therefore, this research is organised as an effort to explore alternative treatment by using EP technique and by focusing on alternative agent to suppress, delay, or reverse carcinogenesis using pharmacologic intervention with naturally occurring or synthetic agents.

1.3 Aim and Objectives

The aim of this research is to investigate the technique and bioactive extract that are able capable to inhibit cancer cell growth (anti-proliferation) by using EP technique and animal extract of PBE test on cultured HeLa cell properties. This study embarks on the following objectives:

- 1) To investigate the suitable parameters of EP technique for Reversible Electroporation (RE) process on HeLa cell.
- 2) To analyze the effect of PBE on HeLa cell lines for anti-proliferation process.
- 3) To evaluate the anti-proliferation mechanism by combining both PBE and the optimized EP method for maximum HeLa cell anti-proliferation activity.

1.4 Scopes of study

In order to achieve the objectives of this research, the following scope of work have been identified such as the process of growing and plating HeLa cell on the surface, expose to pulse electric field (PEF), and to test the PBE efficiency:

1. HeLa cell was cultured in an environment similar to that of the host tissue. The cell were sterile, controlled and maintained at temperature of 37°C and humidified environment at 5% carbon dioxide (CO₂) in order to maintain

the pH, the scale used to specify how acidic or basic a water-based solution of the medium similar to pH of the human blood plasma.

2. Investigation of the best EP parameter for the inhibition growth of HeLa cell in-vitro. EP parameters used:
 - i. Electric field : 100 V/cm – 500 V/cm followed by Standard Electric Field Parameters for Electrochemotherapy (SEFPECT)
 - ii. Pulse Duration : 100 μ s – 5 ms.
 - iii. No of pulses: Single pulse
3. Observation and examination of cellular behavior of HeLa cell in the presence of PBE with exploration of lower extract concentration ranging from 10.0 μ g/ml – 160.0 μ g/ml. Moreover, half maximal inhibitory concentration, IC_{50} was measured to indicate how much concentration of PBE extract (inhibitor) is needed to inhibit the proliferation rate percentage of HeLa cell by half. Therefore, IC_{50} represents the concentration of PBE that is required for 50% inhibition in-vitro of HeLa cell's proliferation rate percentage.
4. The experiments were performed thrice for consistency, and the mean or average proliferation factor was then measured.

1.5 Thesis structure:

The rest of the thesis is structured into eight chapters as follows:

Chapter 2: This chapter covers a background and literature review on EP method. In addition, a brief description of HeLa cell and its natural behaviour/ characteristic are explained in this chapter. Moreover, general information regarding PBE and its benefits use in anti-proliferation application are also discussed in this chapter.

Chapter 3: This chapter provides a list of general materials and equipment used in carrying out the research. Detailed description of the general methods used in achieving the research aim is discussed in this chapter.

Chapter 4: In this chapter, the influence of EP method induced on HeLa cell line and the morphological changes of the cell is investigated, observed and reported.

Chapter 5: In this chapter, the most suitable electric field parameter for the growth of HeLa cell is investigated. The suitability of EP parameters is crucial and is applied in the subsequent chapters to study the influence of EP method on cellular behaviours of the cell and PBE.

Chapter 6: This chapter investigates the effect of PBE on the cellular behaviour of HeLa cell such as the anti-proliferation of the HeLa cell.

Chapter 7: In this chapter, the combined influence of EP and PBE on HeLa cell proliferation rate percentage towards anti-proliferation application are investigated and described.

Chapter 8: This final concludes the thesis and includes recommendations for future work.



CHAPTER 2

LITERATURE REVIEW

2.1 The Cell

Cells are the building blocks of life and are the smallest structural unit and functional organism of all living things (Rodamporn, 2011). The cells are derived from two types of pre-existing cell and managed to accomplish all the activities of cell. In addition, the cells are categorized into two groups, prokaryotic and eukaryotic cell (Rhoads, 2007). An example of a prokaryotic cell is a bacteria cell as shown in Figure 2.1. Eukaryotic cells include plant, animal, fungi and protist (Maskarinec *et al.*, 2006). Contrarily, the size of eukaryotic cells in Figure 2.2 is bigger compared with prokaryotic cells which are smaller in size and it have no nucleus. The cell wall membrane is very flexible with a thin layer surrounding the cell. The cell wall has a thick layer outside the cell membrane and that makes it very tough. The function of this cell wall is to give a physical rigidity and also will allow cellular material and chemical substance to pass through it.




Figure 2.1: A diagram of a Prokaryotic cell
(<https://commons.wikimedia.org>)




Figure 2.2: Eukaryotic cell. Adapted from
(<http://ib.bioninja.com>)

2.2 Composition of cell

2.2.1 Nucleus

One of the most important organelles in a cell is nucleus. Nucleus found in eukaryotic only cells and has two main functions: it retains and stores the cell's hereditary information or genetic information through (DNA); and coordinates and controls cell's activities, which include intermediary metabolism, cell division, protein synthesis, and also cell growth (Will W. Minuth and Raimund Strehl, 2005). The nucleus is composed of a double membrane nuclear envelope that carries the entire organelle sheathed, and the nucleoskeleton that helps in supporting the cell as a whole. The nucleus also controls the safety of the genes and command the activity of the entire cell by managing the gene expression. Thus, the nucleus is referred as the control centre or also known as the brain of the cell (Will W. Minuth and Raimund Strehl, 2005). Figure 2.3 shows the image of cell nucleus.




Figure 2.3: Illustration of Nucleus. Adapted from
(<https://biology.tutorvista.com>)

2.2.2 Cell membrane

Cell membrane or also known as plasma membrane (Figure 2.4) acts as a semi-porous barrier to the external environment. The cell membrane is not only the boundary of the cell unit, but it also has a specific compartment that accommodates many essential cell functions, carrying molecules in and out of cell and helps in metabolic functions (Lombard, J., 2014). In addition, cell membrane allows the cell to communicate with its external surroundings. Hydrophobic tails and hydrophilic heads are the composition of the cell membrane known as phospholipid bilayer. Other than phospholipids, cholesterol are also present in the cell membrane. Cell membrane only allows certain substances to diffuse into it like oxygen (O_2), carbon dioxide (CO_2) and water (H_2O) while ions and some polar molecules can't diffuse through the bilayer because the membrane are made up from lipids due to existence of stability and flexibility in lipids bilayers.




Figure 2.4: Illustration of Cell Membrane. Adapted from
(<http://anatomysciences.com>)

REFERENCES

- Adams, D.J. (2012). *The Valley of Death in Anticancer Drug Development: A Reassessment. Trends in Pharmacological Sciences*. 33, 173-180.
- Adon, M. N. (2015). *Pulsed Electric Field Exposure Effect on Morphological properties of HeLa cell*. Thesis Ph. D. Universiti Tun Hussein Onn Malaysia.
- Ahmed M, Brace CL, Lee FT, Goldberg SN.(2011). *Principles of and advances in percutaneous ablation. Radiology*. 258:351–369.
- Al Ünyayar, Murat Demirbilek, Melisa Turkoglu, Mehmet A. Mazmanci, Emrah A. Erkurt, Serpil Ünyayar, Özlem Cekic, Ayla Celik, and Hatice Atacag. (2005). *Cytotoxic Activities of Funalia trogii (Berk.) Bond. Et. Singer ATCC 200800 Bioactive Extract on HeLa Cell and Fibroblast Cells. International Journal of Medicinal Mushrooms*. Vol. 7, pp. 478-479.
- Anelli T, Sitia R. (2008). *Protein quality control in the early secretory pathway. EMBO J*. 27:315–27.
- Anita, G., Mir, L. M., & Gehl, J. (2003). *Electrochemotherapy: results of cancer treatment using enhanced delivery of bleomycin by electroporation. Cancer treatment reviews, anti-tumour treatment*. 23:371-387.
- Bailón-Moscoso, N., Romero-Benavides, J.C. and Ostrosky-Wegman, P. (2014). *Development of Anticancer Drugs Based on the Hallmarks of Tumor Cells. Tumor Biology*. 35, 3981-3995.
- Barroso, M. D. S. (2014). *The Bezoar Stone: A Princely Antidote, The Távara Sequeira Pinto Collection – Oporto. Acta Medico-Historica Adriatica*. 12(1), 77–98.
- Benedek GB, Villars FMH. (2000). *Physics with illustrative examples from medicine and biology: electricity and magnetism (second edition)*. Springer.
- Boschberg P. (2006). *The trade, forgery and medicinal use of porcupine bezoars in the early modern period (c.1500-1750)*. In: Pinto CA, editor. Oriente. Lisbon: Fundação Oriente, Orient Foundation, 60-78.

- Davalos, R. V. Mir, L. M. & Rubinsky, B. (2005). *Tissue ablation with irreversible electroporation*. Ann Biomed Eng, 33: 223-231.
- Duffin, C. J. (2013). *Porcupine Stones*. *Pharmaceutical Historian*. 43(1).
- Enos Tangke Arung, Britanto Dani Wicaksono, Yohana, Ayupriyanti Handoko, Irawan Wijaya Kusuma, Dina Yulia, and Ferry Sandra. (2009). *Anti-Cancer Properties of Diethylether Extract of Wood from Sukun (Artocarpus altilis) in Human Breast Cancer (T47D)*. 8 (4): pp 317-323.
- Faurie, C., Golzio, M., Phez, E., Teissie, J. & Rols, M. P. (2005). *Electric Field-Induced Cell Membrane Permeabilization and Gene Transfer: Theory and Experiments*. *Engineering in Life Sciences*. 5(2), pp. 179-186.
- Firus Khan, A. Y., Abdullah Asuhaimi, F., Jalal, T., Natto, H. A., Roheem, F. O., Ahmed, Q. U., Johan, M. F., Hashim, R., & Abdul Wahab, R. (2017a). *Anti-Proliferative Property of Porcupine Bezoar Extract on Melanoma (A375)*. *International Journal of Allied Health Sciences*, 1(2), 1-13.
- Firus K AY, Mohd N MH, Tara J, Razali S, Hassan H, Asuhaimi FA, Wahab RA. (2017b). *In Vitro Evaluation of Porcupine Bezoar Extracts as Anticancer Agent on A549 -A Preliminary Study*. *Adv Biotech & Micro*. 5(1): 555-651
- Friedman, J. R., & Nunnari, J. (2014). *Mitochondrial form and function*. *by nature*. 505(7483), 335-343.
- Funian Xiao, Leary, J. F., Reece, L., Camarillo, I. G., Sundararajan, M., & Sundararajan, R. (2010). *Nanosecond, low voltage electro-endocrine-therapy for breast cancer treatment*. Annual Report Conference on Electrical Insulation and Dielectric Phenomena.
- Funian Xiao, Therese Salameh, Ignacio Camarillo, Carly Dykstra, Nick Lenarduzzi, Lisa Reece, Jim F. Leary, Arutselvan Natarajan, and Raji Sundararajan . (2009). *Electrically-Enhanced Chemodrug Delivery to Human Breast Cancer Cells*. Conf Presentation Biomedical.
- Gehl, J. (2003). *Electroporation: theory and methods, perspectives for drug delivery, gene therapy and research*. *Acta Physiologica Scandinavica* 177(4): 437-447.
- HB Mamman, MMA Jamil, MN Adon. (2016). *Optimization of electric field parameters for HT29 cell line towards wound healing application*. *Indian Journal of Science and Technology*.
- Heller, L. C. & Heller, R. (2006). *In vivo electroporation for gene therapy*. *Human Gene Therapy*. 17 (9), pp. 890-897.

- Hjouj, M., D. Last, et al. (2012). *MRI study on reversible and irreversible electroporation induced blood brain barrier disruption*. PLoS One 7(8): e42817.
- Hondroulis, E., Melnick, S. J., Zhang, X., Wu, Z.-Z., & Li, C.-Z. (2013). *Electrical field manipulation of cancer cell behavior monitored by whole cell biosensing device*. Biomedical Microdevices, 15(4), 657–663.
- Huang S., Ingber D.E. (1999). *The structural and mechanical complexity of cell-growth control*. Nat. Cell Biol.; 1:E131.
- Ivorra, A. & Rubinsky, B. (2006). *Impedance analyser for in vivo electroporation studies*. Proceedings of the 28th EMBS Annual International Conference, New York City, USA.
- Jaeger, H., Balasa, A., & Knorr, D. (2009). *Food Industry Applications for Pulsed Electric Fields*. In *Electrotechnologies for Extraction from Food Plants and Biomaterials*. New York: Springer. pp. 181-216.
- Jorgensen, P. & Tyers, M. (2004). *How cells coordinate growth and division*. Current Biology. 14(23), pp. R1014-R1027.
- Kanthou, C., Kranjc, S., Sersa, G., Tozer, G., Zupanic, A., & Cemazar, M. (2006). *The endothelial cytoskeleton as a target of electroporation-based therapies*. Molecular Cancer Therapeutics, 5(12), 3145–3152.
- Kerr JF, Wyllie AH, Currie AR. (1972). *Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics*. Br. J. Cancer.26, 239.
- Korohoda, W., Grys, M., & Madeja, Z. (2013). *Reversible and irreversible electroporation of cell suspensions flowing through a localized DC electric field*. Cellular and Molecular Biology Letters, 18(1).
- Landesberg, R., Cozin, M., Cremers, S., Woo, V., Kousteni, S., Sinha, S. & Raghavan, S. (2008). *Inhibition of oral mucosal cell wound healing by bisphosphonates*. Journal of Oral and Maxillofacial Surgery, 66(5), pp. 839-847.
- Lebar, A. M., Sersa, G. R. E. G. O. R., Kranjc, S. I. M. O. N. A., Groselj, A. & Miklavcic, D. (2002). *Optimisation of pulse parameters in vitro for in vivo electrochemotherapy*. Anticancer Research, 22(3), pp. 731-1736.
- Lloyd, A. C. (2013). *The regulation of cell size*. 154(6), pp. 1194-1205.
- Lombard, J. (2014). *Once upon a time the cell membranes: 175 years of cell boundary research*. Biology Direct, 9, 32.

- M. Heish, T. Salameh, I. Camarillo and R. Sundararajan. (2011). *Irreversible Electroporation Effects: A Drug-Free Treatment for Cancer*.
- Manabe, Y., Nakagawa, R., Zhehong, S., Maetani, M., Teranishi, K., Shimomura, N., & Takahashi, A. (2011). *Influences of pulsed electric fields on the gene expression of pathogenic bacteria*. In Pulsed Power Conference (PPC). IEEE. 1242-1246.
- Marguerat, S., & Bähler, J. (2012). *Coordinating genome expression with cell size*. Trends in Genetics, 28(11), pp. 560-565.
- Martin, A. S. & Richard, K. A. (2001). *Intergrin and cell proliferation: regulation of cyclin-dependent kinases via cytoplasmic signalling pathways*. Journal of cell science 114, Signal transduction and cellular organization, 2553-2560.
- Maskarinec, G. (2006). *A Longitudinal Investigation of Mammographic Density: The Multiethnic Cohort*. Cancer Epidemiology Biomarkers & Prevention, 15(4), 732-739.
- Mehta, K., Gandhi, V., Pathak, S., Aggarwal, B.B. and Grover, R.K. (2014). *Multi-Targeted Approach to Cancer Treatment*. An International Translational Cancer Research Symposium. Anticancer Research, 34, 6791-6795.
- Meier P, Finch A and Evan G. (2000). *Apoptosis in development*. Nature 407: 796 – 801.
- Méthot, S., V. Moulin, et al. (2001). *Morphological changes of human skin cells exposed to a DC electric field in vitro using a new exposure system*. The Canadian Journal of Chemical Engineering 79(4): 668-677.
- Mir, L. M., Orlowski, S., Belehradek, J. & Paoletti, C. (1991). *Electrochemotherapy potentiation of antitumour effect of bleomycin by local electric pulses*. European Journal of Cancer and Clinical Oncology, 27(1), pp. 68-72.
- Miyata, S., Koshikawa, N., Yasumitsu, H. & Miyazaki, K. (2000). *Trypsin stimulates integrin $\alpha 5 \beta 1$ -dependent adhesion to fibronectin and proliferation of human gastric carcinoma cells through activation of proteinase-activated receptor-2*. Journal of Biological Chemistry, 275(7), pp. 4592-4598.
- Moffat, J.G., Rudolph, J. and Bailey, D. (2014). *Phenotypic Screening in Cancer Drug Discovery—Past, Present and Future*. Nature Reviews Drug Discovery, 13, 588-602.
- Mori E, Sforzi A. (2013). *Structure of phytobezoars found in the stomach of a crested porcupine, Hystrix cristata L*. Folia Zool 2013; 62: 232-4.

- Nikon Corporation. (2015). *Inverted Microscope Eclipse TS100 Manual*. Japan: Nikon.
- Oakes, S. A., & Papa, F. R. (2015). *The Role of Endoplasmic Reticulum Stress in Human Pathology*. Annual Review of Pathology, 10, 173–194
- Paula, A. G., John, H. R. & Rafael, V. D. (2011). *Electrical conductivity changes during irreversible electroporation treatment of brain cancer*. Annual International Conference of Engineering in Medicine and Biology Society, EMBC, IEEE, 30 Aug.-3 Sept.
- Pavšelj, N. & Miklavčič, D. (2008). *Numerical modeling in electroporation-based biomedical applications*. Radiology and Oncology, 42(3), pp. 159-168.
- Pehlivanova, V. N., Tsoneva, I. H. & Tzoneva, R. D. (2012). *Multiple effects of electroporation on the adhesive behaviour of breast cancer cells and fibroblasts*. Cancer Cell International, 12(9), pp.1-14.
- Pehlivanova, V., Tsoneva, I. & Tzoneva, R. (2010). *Influence of electroporation on cell adhesion, growth and viability of cancer cells and fibroblasts*. Biologie Cellulaire, 64(4), pp. 581-90.
- Prindull, G. (1995). *Apoptosis in the embryo and tumorigenesis*. European Journal of Cancer, 31(1), 116–123.
- R. Sundararajan (2014). *Electroporation-Based Therapies for Cancer: From Basics to Clinical Applications*. Elsevier, Aug 28, Medical, 352.
- R. Sundararajan, Ramyan Rajendran, Sajan S. Shahid, Santosh D.K, Snehalatha Radhakrishnan, Priyadarshan K., Varsha S., U. Vimal Kumar, Rajaprabu Ramachandran, and Kavita Sankaranarayanan. (2011). *Effect of Irreversible Electroporation on Cancer Cells*. Electrical Insulation and Dielectric Phenomena (CEIDP), Annual Report Conference. IEEE. 16-19 Oct. 2011. 164 – 167.
- R. Sundararajan, T. Salameh, I. Camarillo and L. Campana. (2011). *Effective Use of Electrical Pulses on Cancer Cells to Control Proliferation*. IEEE.
- Rajendran N and Ramakrishnan J. (2010). *Polyphenol analysis and antitumor activity of crude extracts from tegmen of Artocarpus heterophyllus*. The Internet Journal of Alternative Medicine. 7 (2): 61-3.
- Riccardi C, Bruno E. (1996). *Food intake of captive porcupines Hystrix cristata (Rodentia, Hystricidae)*. Atti Soc Tosc Sci Nat Mem B. 103:81-3

- Ridzwan BH, Rosli R, Nihayah M & Asma A. (2003). *Preliminary Study On The Anatomy, Growth and Haematology Parameters of Landak Borneo (Thecurus Crassispinas)*. Proc. 25th Malaysian Soc. Animan Production Annual Conference. 1-3 August, Melaka.
- Rhoads, D. (2007). *History of Cell Biology*.
- Rodamporn, S. (2010). *Microfluidic Systems for Cell Transfection using Sonoporation and Electroporation*. University of Southampton: Ph.D. Thesis.
- Rubinsky, B. (2007). *Irreversible electroporation in medicine*. Technology in Cancer Research & Treatment, 6(4), pp. 255-259.
- Saulis, G. (2012). *Electrochemical Processes Occuring during Cell Electromanipulation Procedures. A short review*. Electroporation in Laboratory and Clinical Investigations, EP Spugnini, A. Baldi (Eds.). pp. 99-113.
- Savill, J., & Fadok, V. (2000). *Corpse clearance defines the meaning of cell death*. Nature, 407(6805), 784-788.
- Serša, G., Štabuc, B., Cemažar, M., Miklavcic, D. & Rudolf, Z. (2000). Electrochemotherapy with cisplatin: the systemic antitumour effectiveness of cisplatin can be potentiated locally by the application of electric pulses in the treatment of malignant melanoma skin metastases. *Melanoma Research*, 10(4), pp. 381-385.
- Skloot, R. (2017). *The immortal life of Henrietta Lacks*. Broadway Books.
- Somphop Rodamporn. (2011). *Optimal Parameters of Electroporation for Gene and Tissue*. The 2011 Biomedical Engineering International Conference (BMEiCON-2011).
- Straughan, D. W., Fentem, J. H. & Balls, M. (1997). *Replacement alternative and complementary in vitro methods in pharmaceutical research*. In vitro methods in pharmaceutical research, academic press, medical, 467.
- Strober, W. (2001). *Trypan blue exclusion test of cell viability*. Current Protocols in Immunology, pp. A3-B.
- Sugar IP, Neumann E. (1984). *Stochastic model for electric field-induced membrane pores electroporation*. BiophysChem 19: 211-225.
- Talele, S., & Gaynor, P. (2008). *Non-linear time domain model of electroporeabilization: Effect of extracellular conductivity and applied electric field parameters*. Journal of Electrostatics, 66(5), pp. 328-334

- Thompson, C. (1995). *Apoptosis in the pathogenesis and treatment of disease*. Science, 267(5203), 1456–1462.
- Titushkin, I. & Cho, M. (2009). *Regulation of cell cytoskeleton and membrane mechanics by electric field: role of linker proteins*. Biophysical Journal, 96(2), pp. 717-728.
- Torre, L. A., Siegel, R. L., Ward, E. M., & Jemal, A. (2015). *Global Cancer Incidence and Mortality Rates and Trends--An Update*. Cancer Epidemiology Biomarkers & Prevention, 25(1), 16–27.
- Tryfona, T., & Bustard, M. T. (2008). *Impact of pulsed electric fields on Corynebacterium glutamicum cell membrane permeabilization*. Journal of Bioscience and Bioengineering, 105(4), pp. 375-382.
- Ünyayar, A., Demirbilek, M., Turkoglu, M., Celik, A., Mazmanci, M. A., Erkurt, E. A., Atacag, H. (2006). *Evaluation of Cytotoxic and Mutagenic Effects of Coriolus versicolor and Funalia trogii Extracts on Mammalian Cells*. Drug and Chemical Toxicology. 29 (1), pp 69–83.
- V. Jothi, V. Raman, R. Sharma, I. Camarillo, and R. Sundararajan. (2016). *Electroporation-based Enhanced Anti-Cancer Effect of Veliparib on Triple Negative Breast Cancer Cells*. Proc. Electrostatics Joint Conference.
- Vinson JA, Su X, Zubik L, Bose P. (2001). *Phenol antioxidant quantity and quality in foods: Fruits*. J Agric Food Chem. 49:5315-21.
- Wang, M., Orwar, O., Olofsson, J. & Weber, S. G. (2010). *Single-cell electroporation. Analytical and Bioanalytical Chemistry*. 397 (8), pp. 3235-3248.
- Will W. Minuth and Raimund Strehl. (2005). *Tissue Engineering: From Cell Biology to Artificial Organs*.
- William J. Dower, Jeff F. Miller and Charles W. Ragsdale. (1988). *High efficiency transformation of E. coli by high voltage electroporation*. Nucleic Acids Research. Volume 16 Number 13.
- Wilson, C., Venditti, R., Rega, L. R., Colanzi, A., D'Angelo, G., & De Matteis, M. A. (2011). *The Golgi apparatus: an organelle with multiple complex functions*. Biochemical Journal, 433(1), 1–9.
- Xiao, F., Otto, K., Camarillo, I., & Sundararajan, R. (2009). *Impedance spectroscopy studies of electroporated human breast cancer cells*. 2009 IEEE Conference on Electrical Insulation and Dielectric Phenomena

- Yang, H. Q., Y. H. Wang, et al. (2012). *Efficacy of Proliferation of HeLa cell under Three Different Low-Intensity Red Lasers Irradiation*. International Journal of Photoenergy 2012: 5.
- Yew PN, Lee WL, Lim YY. (2017). *Antioxidant and intracellular reactive oxygen species/reactive nitrogen species scavenging activities of three porcupine bezoars from Hystrix brachyura*. Phcog Res 2017; 9: 366-71
- Zimmerman U, Pilwat G, Reimann F. (1974). *Dielectric breakdown of cell membranes*. Biophys J14: 881-899.
- Županič, A. & Miklavčič, D. (2010). *Optimization and Numerical Modeling in Irreversible Electroporation Treatment Planning. In Irreversible Electroporation*. Berlin Heidelberg: Springer. pp. 20

